

CLASS – 11

WORKSHEET- WORK, ENERGY, POWER

(1 mark questions)

1. Can a body have energy without momentum?

2. Can a body have momentum without energy?

3. A light body and a heavy body have the same momentum. Which one will have greater kinetic energy?

4. A spark is produced, when two stones are struck against each other. Why?

5. Define power and its SI unit.

6. How many watts are there in one horse power?

7. Give three examples of forces which are conservative in nature.

8. An artificial satellite is at a height of 36,500km above earth's surface. What is the work done by earth's gravitational force in keeping it in its orbit?

9. State work energy theorem.

10. The work energy theorem states that the change in

- (a) kinetic energy of a particle is equal to the work done on it by the net force.
- (b) kinetic energy of a particle is equal to the work done by one of the forces acting on it.
- (c) potential energy of a particle is equal to the work done on it by the net force.
- (d) potential energy of a particle is equal to the work done by one of the forces acting on it.

11. Prove work-energy theorem.

12. A body is being raised to a height h from the surface of earth. What is the sign of work done by (i) applied force (ii) gravitational force?

(a) Positive, positive (b) Positive, Negative (c) Negative, Positive (d) Negative, negative

13. A weightlifter lifts a weight off the ground and holds it up

(a) work is done in lifting as well as holding the weight.


(b) no work is done in both lifting and holding the weight.

(c) work is done in lifting the weight but no work is required to be done in holding it up.

(d) no work is done in lifting the weight but work is required to be done in holding it up.

14. How will the momentum of a body change if its kinetic energy is doubled?

15. A force F acting on an object varies with distance x as shown in the figure. The work done by the force in moving the object from $x = 0$ to $x = 20\text{m}$ is



- (a) 500J (b) 1000J (c) 1500J (d) 2000J

16. The potential energy of a system increases if work is done

- (a) upon the system by a non conservative force
(b) by the system against a conservative force
(c) by the system against a non-conservative force
(d) upon the system by a conservative force

17. The negative of the work done by the conservative internal forces on a system equals to the change in

- (a) total energy (b) kinetic energy (c) potential energy (d) none of these

18. Can kinetic energy be negative? What about potential energy?

19. A body is falling freely under the action of gravity alone in vacuum. Which of the following quantities remain constant during the fall?

- (a) Kinetic energy (b) Potential energy
(c) Total mechanical energy (d) Total linear momentum

20. Two springs of spring constants 1000N m^{-1} and 2000N m^{-1} are stretched with same force. They will have potential energy in the ratio of
(a) 2:1 (b) $2^2:1^2$ (c) 1:2 (d) $1^2:2^2$

21. The power of a water pump is 2kW. If $g = 10\text{ ms}^{-2}$, the amount of water it can raise in one minute to a height of 10m is
(a) 2000 litre (b) 1000 litre (c) 100 litre (d) 1200 litre

22. Water is flowing in a river at 2m/s. The river is 50m wide and has an average depth of 5m. The power available from the current in the river is (density of water = 1000 kg m^{-3})
(a) 0.5 MW (b) 1 MW (c) 1.5 MW (d) 2 MW

23. What is the value of coefficient of restitution in (i) perfectly elastic collision and (ii) perfectly inelastic collision?

24. The sign of work done by a force on a body is important to understand. State carefully if the following quantities are positive or negative: (1 mark each)

(a) Work done by a man in lifting a bucket out of a well by means of a rope tied to the bucket,

(b) Work done by gravitational force in the above case,

(c) Work done by friction on a body sliding down an inclined plane,

(d) Work done by an applied force on a body moving on a rough horizontal plane with uniform velocity,

(e) Work done by the resistive force of air on a vibrating pendulum in bringing it to rest.

25. Point out the correct alternative:

(a) When a conservative force does positive work on a body, the potential energy of the body increases/decreases/remains unaltered.

(b) Work done by a body against friction always results in a loss of its kinetic/potential energy.

(c) The rate of change of total momentum of a many-particle system is proportional to the external force/sum of the internal forces of the system.

(d) In an inelastic collision of two bodies, the quantities which do not change after the collision are the total kinetic energy/total linear momentum/total energy of the system of two bodies.

26. State if each of the following statements is true or false. Give reasons for your answer.

(a) In an elastic collision of two bodies, the momentum and energy of each body is conserved.

(b) Total energy of a system is always conserved, no matter what internal and external forces on the body are present.

(c) Work done in the motion of a body over a closed loop is zero for every force in nature.


(d) In an inelastic collision, the final kinetic energy is always less than the initial kinetic energy of the system.

27. A body is initially at rest. It undergoes a one-dimensional motion with constant acceleration. The power delivered to it at time t is proportional to

(i) $t^{1/2}$ (ii) t (iii) $t^{3/2}$ (iv) t^2

28. A body is moving unidirectionally under the influence of a source of constant power. Its displacement in time t is proportional to
 (i) $t^{1/2}$ (ii) t (iii) $t^{3/2}$ (iv) t^2

29. Two identical ball bearings in contact with each other and resting on a friction less table are hit head-on by another ball bearing of the same mass moving initially with a speed V . If the collision is elastic, which of the following (Fig.) is a possible result after collision?



(2 marks Qs)

30. A force $\vec{F} = \hat{i} + 5\hat{j} + 7\hat{k}$ acts on a particle and displaces it through $\vec{s} = 6\hat{i} + 9\hat{k}$. Calculate the work done if the force is in newton and displacement in metre. [Ans. 69J]

31. How high must a body be lifted to gain an amount of potential energy equal to the kinetic energy it has when moving at speed 20m/s? The value of acceleration due to gravity at a place is $g = 9.8 \text{ m/s}^2$. [Ans. 20.2m]

32. Calculate the work done in lifting a 300N weight to a height of 10m with an acceleration 0.5 ms^{-2} . Take $g = 10 \text{ ms}^{-2}$. [Ans. 3150J]

33. A man weighs 60 kg climbs up a staircase carrying a load of 20kg on his head. The stair case has 20 steps each of height 0.2m. If he takes 10s to climb, find his power.

[Ans. 313.6W]

34. Two bodies of masses m_1 and m_2 have the same linear momentum. What is the ratio of their kinetic energies?

35. Can there be a solution in which $E - U < 0$?

36. An elastic spring of force constant k is compressed by an amount x . Show that its potential energy is $\frac{1}{2} kx^2$.

37. Define work, power and energy and give their SI units.

-
-
38. If a force $\vec{F} = (-2\hat{i} + 3\hat{j} + \hat{k})$, causes a displacement $\vec{S} = (\hat{i} + 2\hat{j} - 4\hat{k})$, of an object, what will be the work done on the object?


39. Find the work done if a particle moves from position $\vec{r}_1 = (4\hat{i} + 3\hat{j} + 6\hat{k})\text{m}$ to a position $\vec{r}_2 = (14\hat{i} + 13\hat{j} + 16\hat{k})\text{m}$ under the effect of force $\vec{F} = (4\hat{i} + 4\hat{j} - 4\hat{k})\text{N}$?

40. A particle is projected making an angle of 45° with the horizontal having kinetic energy K . What is the kinetic energy of highest point?

41. A ball is dropped vertically from rest at a height of 12m. After striking the ground it bounces at a height of 1m. What fraction of kinetic energy does it lose on striking the ground?

42. Given figures are examples of some potential energy functions in one dimension. The total energy of the particle is indicated by a cross on the ordinate axis. In each case,

specify the regions, if any, in which the particle cannot be found for the given energy. Also, indicate the minimum total energy the particle must have in each case. Think of some physical contexts for which these potential energy shapes are relevant.



43. A particle of mass 0.5 kg travels in a straight line with velocity $u = a x^{3/2}$, where $a = 5 \text{ m}^{1/2} \text{ s}^{-1}$. What is the work done by the net force during its displacement from $x = 0$ to $x = 2 \text{ m}$? [Ans. 50J]

(3 marks Qs)

44. A man weighing 50 kg supports a body of 25 kg on his head. What is the work done when he moves a distance of 20m up in an incline of 1 in 10? Take $g = 9.8 \text{ ms}^{-2}$. [Ans. 1470J]

45. A shot travelling at the rate of 100 ms^{-1} is just able to pierce a plank 4cm thick. What velocity is required to just pierce a plank 9cm thick? [Ans. 150m/s]

46. If the kinetic energy of a body increases by 300%, by what % will the linear momentum of the body increase? [Ans. 100%]

47. While catching a cricket ball of mass 200g moving with a velocity of 20m/s, the player draws his hands backwards through 20cm. Find the work done in catching the ball and the average force exerted by the ball on the hand. [Ans. 40J, 200N]

48. Two springs have force constants k_1 and k_2 ($k_1 > k_2$). On which spring is more work done, if (i) they are stretched by the same factor and (ii) they are stretched by the same amount?

49. A block of mass 2kg is dropped from a height of 40cm on a spring whose force-constant is 1960 Nm^{-1} . What will be the maximum distance x through which the spring is compressed? [Ans. 10cm]

50. The human heart discharges 75ml of blood at each beat against a pressure of 0.1m of Hg. Calculate power of heart assuming that pulse frequency is 80 beats per minute. Density of Hg = $13.6 \times 10^3 \text{ kgm}^{-3}$. [Ans. 1.33W]

51. A man cycles up a hill, whose slope is 1m in 20 with it velocity of 6.4 kmh^{-1} . The weight of man and the cycle is 98kg. What work per minute is he doing? What is his horse power? [Ans, 0.144 hp, 5122.1J]

52. A 10kg ball and 20kg ball approach each other with velocities 20 ms^{-1} and 10 ms^{-1} respectively. What are their velocities after collision if the collision is perfectly elastic?
[Ans. 20 ms^{-1} , 10 ms^{-1}]

53. A ball of 0.1kg makes an elastic head on collision with a ball of unknown mass that is initially at rest. If the 0.1kg ball rebounds at one-third of its original speed, what is the mass of the other ball?
[Ans. 0.2 kg]

54. Define the term potential energy, and derive its dimensions. Write an expression for the gravitational potential energy of a body of mass m raised to a height h above the earth's surface.


55. What is the meaning of 'Collision' in physics? Differentiate between elastic and inelastic collision. Give one example each.

56. Show that the total mechanical energy of a body falling freely under gravity is conserved. Discuss it graphically.

57. Show that in case of one dimensional elastic collision of two bodies, the relative velocity of separation after the collision is equal to the relative velocity of approach before the collision.

58. Prove that bodies of identical masses exchange their velocities after head on elastic collision.

59. Figure gives the acceleration of a 2.0kg body as it moves from rest along x-axis while a valuable force acts on it from $x = 0\text{m}$ to $x = 9\text{m}$. Find the work done by the force on the body when it reaches (i) $x = 4\text{m}$ and (ii) $x = 7\text{m}$.




60. A bullet of mass 0.012kg and horizontal speed 70 m/s strikes a block of wood of mass 0.4kg and instantly comes to rest with respect to the block. The block is suspended from the ceiling by means of thin wires. Calculate the height to which the block rises. Also, estimate the amount of heat produced in the block.

61. If the elongation in a spring of force constant k is tripled, calculate
 (a) ratio of final to initial force in the spring.
 (b) ratio of elastic energies stored in the two cases
 (c) work done in changing to the state of elongation.


62. The spring shown in figure has a force constant 24 n/m . The mass of the block attached to the spring is 4kg . Initially the block is at rest and spring is unstretched. The horizontal

force of 10N is applied on the block, then what is the speed of the block when it has been moved through a distance of 0.5m?



63. The turbine pits at the Niagra falls are 50m deep. The average horse power developed is 5000. If the efficiency of the generator is 85%, how much water passes through the turbines per minute? Take $g = 10 \text{ ms}^{-2}$.

64. The potential energy function for a particle executing linear simple harmonic motion is given by $V(x) = kx^2/2$, where k is the force constant of the oscillator. For $k = 0.5 \text{ Nm}^{-1}$, the graph of $V(x)$ versus x is shown in Fig. Show that a particle of total energy 1 J moving under this potential must 'turn back' when it reaches $x = \pm 2 \text{ m}$.



-
-
65. A body constrained to move along the z-axis of a coordinate system is subject to a constant force F given by

$$\vec{F} = -\hat{i} + 2\hat{j} + 3\hat{k} \text{ N}$$

where \hat{i} , \hat{j} and \hat{k} are unit vectors along the x- y- and z-axis of the system respectively.

What is the work done by this force in moving the body a distance of 4 m along the z-axis?
[Ans. 12J]


-
-
-
-
-
-
66. An electron and a proton are detected in a cosmic ray experiment, the first with kinetic energy 10 keV, and the second with 100 keV. Which is faster, the electron or the proton? Obtain the ratio of their speeds, (electron mass = 9.11×10^{-31} kg, proton mass = 1.67×10^{-27} kg, $1 \text{ eV} = 1.60 \times 10^{-19} \text{ J}$).
[Ans. 13.53]

-
-
-
-
-
-
67. A raindrop of radius 2 mm falls from a height of 500 m above the ground. It falls with decreasing acceleration (due to viscous resistance of the air) until at half its original height, it attains its maximum (terminal) speed, and moves with uniform speed thereafter. What is the work done by the gravitational force on the drop in the first and second half of its journey? What is the work done by the resistive force in the entire journey if its speed on reaching the ground is 10 ms^{-1} ?
[Ans. 0.1623 J]
-
-
-
-
-
-

68. A molecule in a gas container hits a horizontal wall with speed 200 ms^{-1} and angle 30° with the normal, and rebounds with the same speed. Is momentum conserved in the collision? Is the collision elastic or inelastic?

69. A pump on the ground floor of a building can pump up water to fill a tank of volume 30 m^3 in 15 min. If the tank is 40 m above the ground, and the efficiency of the pump is 30%, how much electric power is consumed by the pump? [Ans. 43.6 kW]

70. The bob A of a pendulum released from 30° to the vertical hits another bob B of the same mass at rest on a table as shown in Fig. How high does the bob A rise after the collision? Neglect the size of the bobs and assume the collision to be elastic.



71. The bob of a pendulum is released from a horizontal position. If the length of the pendulum is 1.5 m, what is the speed with which the bob arrives at the lowermost point, given that it dissipated 5% of its initial energy against air resistance? [Ans. 5.3 m/s]


72. A trolley of mass 300 kg carrying a sandbag of 25 kg is moving uniformly with a speed of 27 km/h on a friction less track. After a while, sand starts leaking out of a hole on the trolley's floor at the rate of 0.05 kg s^{-1} . What is the speed of the trolley after the entire sand bag is empty?

73. A person trying to lose weight (dieter) lifts a 10 kg mass, one thousand times, to a height of 0.5 m each time. Assume that the potential energy lost each time she lowers the mass is dissipated, (a) How much work does she do against the gravitational force? (b) Fat supplies $3.8 \times 10^7 \text{ J}$ of energy per kilogram which is converted to mechanical energy with a 20% efficiency rate. How much fat will the dieter use up? [Ans. 49000J, $6.45 \times 10^{-3} \text{ kg}$]

74. A family uses 8 kW of power, (a) Direct solar energy is incident on the horizontal surface at an average rate of 200 W per square meter. If 20% of this energy can be converted to useful electrical energy, how large an area is needed to supply 8 kW? (b) Compare this area to that of the roof of a typical house. [Ans. 200 m^2 , 21:21]


75. A bullet of mass 0.012 kg and horizontal speed 70 ms^{-1} strikes a block of wood of mass 0.4 kg and instantly comes to rest with respect to the block. The block is suspended from the ceiling by thin wire. Calculate the height to which the block rises. Also, estimate the amount of heat produced in the block. [Ans. 0.212m , 28.54J]

76. Two inclined frictionless tracks, one gradual and the other steep meet at A from where two stones are allowed to slide down from rest, one on each track (Fig). Will the stones reach the bottom at the same time? Will they reach there at the same speed? Explain. Given $\theta_1 = 30^\circ$, $\theta_2 = 60^\circ$, and $h = 10 \text{ m}$, what are the speeds and times taken by the two stones?



[Ans. 14.14 ms^{-1}]

77. A 1 kg block situated on a rough incline is connected to a spring with spring constant 100 Nm^{-1} as shown in Figure. The block is released from rest with the spring in the unstretched position. The block moves 10 cm down the incline before coming to rest. Find the coefficient of friction between the block and the incline. Assume that the spring has negligible mass and the pulley is frictionless.




[Ans. 0.126]


78. A bolt of mass 0.3 kg falls from the ceiling of an elevator moving down with a uniform speed of 7 ms^{-1} . It hits the floor of the elevator (length of elevator = 3 m) and does not rebound. What is the heat produced by the impact? Would your answer be different if the elevator were stationary? [Ans. 8.82 J]

79. A trolley of mass 200 kg moves with a uniform speed of 36 km h^{-1} on a friction less track. A child of mass 20 kg runs on the trolley from one end to the other (10 m away) with a speed of 4 ms^{-1} relative to the trolley in a direction opposite to the trolley's motion, and jumps out of the trolley. What is the final speed of the trolley? How much has the trolley moved from the time the child begins to run? [Ans. 10.36 ms^{-1} , 25.9 m]

80. Which of the following potential energy curves in Fig. cannot possibly describe the elastic collision of two billiard balls? Here r is distance between centres of the balls.



81. Consider the decay of a free neutron at rest: $n \rightarrow p + e^-$. Show that the two body decay of this type must necessarily give an electron of fixed energy, and therefore, cannot account for the observed continuous energy distribution in the β -decay of a neutron or a nucleus, Fig.



85. A body of mass 2 kg initially at rest moves under the action of an applied horizontal force of 7 N on a table with coefficient of kinetic friction = 0.1. Compute the
- (a) Work done by the applied force in 10 s
 - (b) Work done by friction in 10 s
 - (c) Work done by the net force on the body in 10 s
 - (d) Change in kinetic energy of the body in 10 s and interpret your results.

[Ans. 882 J, - 246.9J, 635J, 635J]

86. Answer the following:
- (a) The casing of a rocket in flight bums up due to friction. At whose expense is the heat energy required for burning obtained? The rocket or the atmosphere?
 - (b) Comets move around the sun in highly elliptical orbits. The gravitational force on the comet due to the sun is not normal to the comet's velocity in general. Yet the work done by the gravitational force over every complete orbit of the comet is zero. Why?
 - (c) An artificial satellite orbiting the earth in very thin atmosphere loses its energy gradually due to dissipation against atmospheric resistance, however small. Why then does its speed increase progressively as it comes closer and closer to the earth?
 - (d) In Fig.(i), the man walks 2 m carrying a mass of 15 kg on his hands. In Fig. (ii), he walks the same distance pulling the rope behind him. The rope goes over a pulley, and a mass of 15 kg hangs at its other end. In which case is the work done greater?

