


Class- X


Mathematics-Basic (241)**Marking Scheme SQP-2020-21**

Max. Marks: 80

Duration:3hrs


1	$156 = 2^2 \times 3 \times 13$	1
2	Quadratic polynomial is given by $x^2 - (a + b)x + ab$ $x^2 - 2x - 8$	1
3	HCF X LCM = product of two numbers $LCM(96, 404) = \frac{96 \times 404}{HCF(96, 404)} = \frac{96 \times 404}{4}$ LCM = 9696 OR Every composite number can be expressed (factorized) as a product of primes, and this factorization is unique, apart from the order in which the factors occur.	$\frac{1}{2}$ $\frac{1}{2}$ 1
4	$x - 2y = 0$ $3x + 4y - 20 = 0$ $\frac{1}{3} \neq \frac{-2}{4}$ As, $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ is one condition for consistency. Therefore, the pair of equations is consistent.	$\frac{1}{2}$ $\frac{1}{2}$
5	1	1
6	$\theta = 60^\circ$ Area of sector = $\frac{\theta}{360^\circ} \pi r^2$ $A = \frac{60^\circ}{360^\circ} \times \frac{22}{7} \times (6)^2 \text{ cm}^2$ $A = \frac{1}{6} \times \frac{22}{7} \times 36 \text{ cm}^2$ $= 18.86 \text{ cm}^2$	$\frac{1}{2}$ $\frac{1}{2}$


	OR Another method- Horse can graze in the field which is a circle of radius 28 cm. So, required perimeter = $2\pi r = 2 \cdot \pi(28)$ cm $= 2 \times \frac{22}{7} \times (28) \text{ cm}$ $= 176 \text{ cm}$	$\frac{1}{2}$ $\frac{1}{2}$
7	By converse of Thale's theorem $DE \parallel BC$ $\angle ADE = \angle ABC = 70^\circ$ Given $\angle BAC = 50^\circ$ $\angle ABC + \angle BAC + \angle BCA = 180^\circ$ (Angle sum prop of triangles) $70^\circ + 50^\circ + \angle BCA = 180^\circ$ $\angle BCA = 180^\circ - 120^\circ = 60^\circ$ OR $EC = AC - AE = (7 - 3.5) \text{ cm} = 3.5 \text{ cm}$ $\frac{AD}{BD} = \frac{2}{3}$ and $\frac{AE}{EC} = \frac{3.5}{3.5} = 1$ So, $\frac{AD}{BD} \neq \frac{AE}{EC}$ Hence, By converse of Thale's Theorem, DE is not Parallel to BC.	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
8	Length of the fence = $\frac{\text{Total cost}}{\text{Rate}}$ $= \frac{\text{Rs.5280}}{\text{Rs 24/metre}} = 220 \text{ m}$ So, length of fence = Circumference of the field $\therefore 220\text{m} = 2 \pi r = 2 \times \frac{22}{7} \times r$ So, $r = \frac{220 \times 7}{2 \times 22} \text{ m} = 35 \text{ m}$	$\frac{1}{2}$ $\frac{1}{2}$
9	 Sol: $\tan 30^\circ = \frac{AB}{BC}$ $\frac{1}{\sqrt{3}} = \frac{AB}{8}$ $AB = 8 / \sqrt{3} \text{ metres}$ Height from where it is broken is $8/\sqrt{3} \text{ metres}$	$\frac{1}{2}$ $\frac{1}{2}$

15	 $\tan 60^\circ = \frac{h}{15}$ $\sqrt{3} = \frac{h}{15}$ $h = 15\sqrt{3} \text{ m}$	$\frac{1}{2}$ $\frac{1}{2}$
16	1	1
17 i)	Ans : b) Cloth material required = $2 \times \text{S A of hemispherical dome}$ $= 2 \times 2\pi r^2$ $= 2 \times 2 \times \frac{22}{7} \times (2.5)^2 \text{ m}^2$ $= 78.57 \text{ m}^2$	1
ii)	a) Volume of a cylindrical pillar = $\pi r^2 h$	1
iii)	b) Lateral surface area = $2 \times 2\pi r h$ $= 4 \times \frac{22}{7} \times 1.4 \times 7 \text{ m}^2$ $= 123.2 \text{ m}^2$	1
iv)	d) Volume of hemisphere = $\frac{2}{3} \pi r^3$ $= \frac{2}{3} \times \frac{22}{7} \times (3.5)^3 \text{ m}^3$ $= 89.83 \text{ m}^3$	1
v)	b) Sum of the volumes of two hemispheres of radius 1cm each = $2 \times \frac{2}{3} \pi 1^3$ Volume of sphere of radius 2cm = $\frac{4}{3} \pi 2^3$ So, required ratio is $\frac{2 \times \frac{2}{3} \pi 1^3}{\frac{4}{3} \pi 2^3} = 1:8$	$\frac{1}{2}$ $\frac{1}{2}$

18 i)	c) (0,0)	1
ii)	a) (4,6)	1
iii)	a) (6,5)	1
iv)	a) (16,0)	1
v)	b) (-12,6)	1
19 i)	c) 90°	1
ii)	b) SAS	1
iii)	b) 4 : 9	1
iv)	d) Converse of Pythagoras theorem	1
v)	a) 48 cm ²	1
20 i)	d) parabola	1
ii)	a) 2	1
iii)	b) -1, 3	1
iv)	c) $x^2 - 2x - 3$	1
v)	d) 0	1
21	Let P(x,y) be the required point. Using section formula $\left\{ \frac{m_1x_2+m_2x_1}{m_1+m_2}, \frac{m_1y_2+m_2y_1}{m_1+m_2} \right\} = (x, y)$ $x = \frac{3(8)+1(4)}{3+1}, \quad y = \frac{3(5)+1(-3)}{3+1}$ $x = 7 \quad y = 3$ (7,3) is the required point	1 1


	OR Let P(x, y) be equidistant from the points A(7,1) and B(3,5) Given AP = BP. So, $AP^2 = BP^2$ $(x-7)^2 + (y-1)^2 = (x-3)^2 + (y-5)^2$ $x^2 - 14x + 49 + y^2 - 2y + 1 = x^2 - 6x + 9 + y^2 - 10y + 25$ $x - y = 2$	1 1
22	By BPT, $\frac{AM}{MB} = \frac{AL}{LC} \dots\dots\dots(1)$ Also, $\frac{AN}{ND} = \frac{AL}{LC} \dots\dots\dots(2)$ By Equating (1) and (2) $\frac{AM}{MB} = \frac{AN}{ND}$	$\frac{1}{2}$ $\frac{1}{2}$ 1
23	To prove: $AB + CD = AD + BC$. Proof: $AS = AP$ (Length of tangents from an external point to a circle are equal) $BQ = BP$ $CQ = CR$ $DS = DR$ $AS + BQ + CQ + DS = AP + BP + CR + DR$ $(AS + DS) + (BQ + CQ) = (AP + BP) + (CR + DR)$ $AD + BC = AB + CD$	1 1
24	For the correct construction	2

25	15 cot A = 8, find sin A and sec A. Cot A = 8/15  $\frac{Adj}{Oppo} = 8/15$ By Pythagoras Theorem $AC^2 = AB^2 + BC^2$ $AC = \sqrt{(8x)^2 + (15x)^2}$ $AC = 17x$ Sin A = 15/17 Cos A = 8/17 OR By Pythagoras Theorem $QR = \sqrt{(13)^2 - (12)^2} \text{ cm}$ $QR = 5\text{cm}$ Tan P = 5/12 Cot R = 5/12 Tan P - Cot R = 5/12 - 5/12 = 0	1 $\frac{1}{2}$ $\frac{1}{2}$ 1 1
26	9, 17, 25, $S_n = 636$ $a = 9$ $d = a_2 - a_1$ $= 17 - 9 = 8$ $S_n = \frac{n}{2} [2a + (n-1)d]$ $S_n = \frac{n}{2} [2a + (n-1)d]$	$\frac{1}{2}$ $\frac{1}{2}$

	$636 = \frac{n}{2} [2 \times 9 + (n-1) 8]$ $1272 = n [18 + 8n - 8]$ $1272 = n [10 + 8n]$ $8n^2 + 10n - 1272 = 0$ $4n^2 + 5n - 636 = 0$ $n = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $n = \frac{-5 \pm \sqrt{5^2 - 4 \times 4 \times (-636)}}{2 \times 4}$ $n = \frac{-5 \pm 101}{8}$ $n = \frac{96}{8} \quad \quad \quad n = \frac{-106}{8}$ $n = 12 \quad \quad \quad n = \frac{-53}{4}$ n=12 (since n cannot be negative)	$\frac{1}{2}$ $\frac{1}{2}$
27	Let $\sqrt{3}$ be a rational number. Then $\sqrt{3} = p/q$ HCF (p,q) = 1 Squaring both sides $(\sqrt{3})^2 = (p/q)^2$ $3 = p^2/q^2$ $3q^2 = p^2$ 3 divides p^2 » 3 divides p 3 is a factor of p Take $p = 3C$ $3q^2 = (3C)^2$ $3q^2 = 9C^2$ 3 divides q^2 » 3 divides q 3 is a factor of q Therefore 3 is a common factor of p and q It is a contradiction to our assumption that p/q is rational. Hence $\sqrt{3}$ is an irrational number.	1 $\frac{1}{2}$ $\frac{1}{2}$ 1
28		

	OR (i) Probability of getting A king of red colour. $P(\text{King of red colour}) = 2/52 = 1/26$ (ii) Probability of getting A spade $P(\text{a spade}) = 13/52 = 1/4$ (iii) Probability of getting The queen of diamonds $P(\text{a the queen of diamonds}) = 1/52$	1 1 1
31	$r_1 = 6\text{cm}$ $r_2 = 8\text{cm}$ $r_3 = 10\text{cm}$ Volume of sphere = $\frac{4}{3}\pi r^3$ Volume of the resulting sphere = Sum of the volumes of the smaller spheres. $\frac{4}{3}\pi r^3 = \frac{4}{3}\pi r_1^3 + \frac{4}{3}\pi r_2^3 + \frac{4}{3}\pi r_3^3$ $\frac{4}{3}\pi r^3 = \frac{4}{3}\pi (r_1^3 + r_2^3 + r_3^3)$ $r^3 = 6^3 + 8^3 + 10^3$ $r^3 = 1728$ $r = \sqrt[3]{1728}$ $r = 12\text{ cm}$ Therefore, the radius of the resulting sphere is 12cm.	1 1 1
32	$(\sin A - \cos A + 1) / (\sin A + \cos A - 1) = 1 / (\sec A - \tan A)$ L.H.S. divide numerator and denominator by $\cos A$ $= (\tan A - 1 + \sec A) / (\tan A + 1 - \sec A)$ $= (\tan A - 1 + \sec A) / (1 - \sec A + \tan A)$ We know that $1 + \tan^2 A = \sec^2 A$ Or $1 = \sec^2 A - \tan^2 A = (\sec A + \tan A)(\sec A - \tan A)$ $= (\sec A + \tan A - 1) / [(\sec A + \tan A)(\sec A - \tan A) - (\sec A - \tan A)]$ $= (\sec A + \tan A - 1) / (\sec A - \tan A)(\sec A + \tan A - 1)$	1 1 1

[illegible]



Let AB and CD be the multi-storeyed building and the building respectively.

Let the height of the multi-storeyed building = h m and
the distance between the two buildings = x m.

$$AE = CD = 8 \text{ m [Given]}$$

$$BE = AB - AE = (h - 8) \text{ m}$$

and

$$AC = DE = x \text{ m [Given]}$$

Also,

$$\angle FBD = \angle BDE = 30^\circ \text{ (Alternate angles)}$$

$$\angle FBC = \angle BCA = 45^\circ \text{ (Alternate angles)}$$

Now,

In ΔACB ,

$$\Rightarrow \tan 45^\circ = \frac{AB}{AC} \left[\because \tan \theta = \frac{\text{Perpendicular}}{\text{Base}} \right]$$

$$\Rightarrow 1 = \frac{h}{x}$$

$$\Rightarrow x = h \dots (i)$$

In ΔBDE ,

1

 $\frac{1}{2}$

1

$$\Rightarrow \tan 30^\circ = \frac{BE}{ED}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h-8}{x}$$

$$\Rightarrow x = \sqrt{3}(h-8) \dots \dots \dots (ii)$$

From (i) and (ii), we get,

$$h = \sqrt{3}h - 8\sqrt{3}$$

$$\sqrt{3}h - h = 8\sqrt{3}$$

$$h(\sqrt{3} - 1) = 8\sqrt{3}$$

$$h = \frac{8\sqrt{3}}{\sqrt{3}-1}$$

$$h = \frac{8\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$$



$$h = 4\sqrt{3}(\sqrt{3}+1)$$

$$h = 12 + 4\sqrt{3} \text{ m}$$

Distance between the two building

$$x = (12 + 4\sqrt{3}) \text{ m} \quad [From(i)]$$

OR



From the figure, the angle of elevation for the first position of the balloon $\angle EAD = 60^\circ$ and for second position $\angle BAC = 30^\circ$. The vertical distance

$$ED = CB = 88.2 - 1.2 = 87 \text{ m}.$$

	Let AD = x m and AB = y m. Then in right $\triangle ADE$, $\tan 60^\circ = \frac{DE}{AD}$ $\sqrt{3} = \frac{87}{x}$ $x = \frac{87}{\sqrt{3}} \dots\dots\dots(i)$ In right $\triangle ABC$, $\tan 30^\circ = \frac{BC}{AB}$ $\frac{1}{\sqrt{3}} = \frac{87}{y}$ $y = 87\sqrt{3} \dots\dots\dots(ii)$ Subtracting(i) and (ii) $y-x = 87\sqrt{3} - \frac{87}{\sqrt{3}}$ $y-x = \frac{87 \cdot 2 \cdot \sqrt{3}}{\sqrt{3} \cdot \sqrt{3}}$ $y-x = 58\sqrt{3} \text{ m}$ Hence, the distance travelled by the balloon is equal to BD $y-x = 58\sqrt{3} \text{ m.}$	1 1 1 1
35	Let A be the first term and D the common difference of A.P. $Tp = a = A + (p-1)D = (A-D) + pD \quad (1)$ $Tq = b = A + (q-1)D = (A-D) + qD \quad \dots(2)$ $Tr = c = A + (r-1)D = (A-D) + rD \quad \dots(3)$ Here we have got two unknowns A and D which are to be eliminated. We multiply (1),(2) and (3) by $q-r$, $r-p$ and $p-q$ respectively and add: $a(q-r) = (A-D)(q-r) + Dp(q-r)$ $b(r-p) = (A-D)(r-p) + Dq(r-p)$ $c(p-q) = (A-D)(p-q) + Dr(p-q)$ $a(q-r) + b(r-p) + c(p-q)$ $= (A-D)[q-r+r-p+p-q] + D[p(q-r) + q(r-p) + r(p-q)]$ $= (A-D)(0) + D[pq-pr+qr-pq+rp-rq]$ $= 0$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 1

36	Height (in cm)	f	C.F.	
	below 140	4	4	
	140-145	7	11	1
	145-150	18	29	
	150-155	11	40	
	155-160	6	46	
	160-165	5	51	
	$N=51 \Rightarrow$			
	$N/2=51/2=25.5$			
	As 29 is just greater than 25.5, therefore median class is 145-150.			
	$Median = l + \frac{(\frac{N}{2} - C)}{f} \times h$			
	Here, l = lower limit of median class = 145			
	C = C.F. of the class preceding the median class = 11			$\frac{1}{2}$
	h = higher limit - lower limit = 150 - 145 = 5			
	f = frequency of median class = 18			
	$\therefore median =$			
	$= 145 + \frac{(25.5 - 11)}{18} \times 5$			$\frac{1}{2}$
	$= 149.03$			
	Mean by direct method			1
	Height (in cm)	f	x_i	fx_i
	below 140	4	137.5	550
	140-145	7	142.5	997.5
	145-150	18	147.5	2655
	150-155	11	152.5	1677.5
	155-160	6	157.5	945
	160-165	5	162.5	812.5
			$\sum fx$	
	Mean	=	$\frac{\quad}{N}$	1
			$= 7637.5/51$	
			$= 149.75$	1